arXiv:0902.0834v1 [hep-ph] 5 Feb 2009

November, 2008

# Neutron-Anti-Neutron Oscillation: Theory and Phenomenology

**R. N. Mohapatra**

*Maryland Center for Fundamental Physics and Department of Physics, University of Maryland, College Park, Maryland, 20742*

**Abstract**

The discovery of neutrino masses has provided strong hints in favor of the possibility that B-L symmetry is an intimate feature of physics beyond the standard model. I discuss how important information about this symmetry as well as other scenarios for TeV scale new physics can be obtained from the baryon number violating process, $n-\bar{n}$ oscillation. This article presents an overview of different aspects of neutron-anti-neutron oscillation and is divided into the following parts : (i) the phenomenon; (ii) the physics, (iii) plausible models and (iv) applications to cosmology. In particular, it is argued how the discovery of $n-\bar{n}$ oscillation can significantly affect our thinking about simple grand unified theory paradigms for physics beyond the standard model, elucidate the nature of forces behind neutrino mass and provide a new microphysical view of the origin of matter in the universe.

# 1 Introduction

Particle oscillations are familiar phenomena in both classical and quantum mechanics and have provided a wealth of information about the nature of matter and forces acting on them. The most elementary example of a pendulum oscillation is caused by the force of gravity and therefore provides information about the strength of gravity. In the domain of quantum mechanics, two states close by in energy can transmute into one another if the Hamiltonian for the system includes a force that connects them and if certain coherence conditions are satisfied. Turning this question around, one can conclude that if experimentally an oscillation between states, forbidden in the standard model is observed, it will then imply the existence of hitherto unknown classes of interactions. Many such instances in the domain of particle physics are known where study of particle oscillations have provided important clarification about the fundamental nature of underlying physics. Kaon and neutrino oscillations are but two glowing examples of this. The first in combination with the recently observed $B - \bar{B}$ oscillations confirmed the nature of CP violation in standard model. The latter has for the first time revealed the existence of new physics as well as new symmetries beyond the standard model.

The question then arises as to whether there are other such possibilities that need to be experimentally explored to probe physics beyond the standard model (SM). In this review, I argue that neutron-anti-neutron ($n - \bar{n}$) oscillation provides one such unique example and is extremely timely in the aftermath of one of the fundamental discoveries in our field i.e. the neutrinos are massive and can provide important clues to our understanding of new symmetries and forces behind the neutrino mass generation.

The $n - \bar{n}$ oscillation[1] is a unique kind of oscillation phenomenon compared to kaon oscillations in that it breaks one of the once sacred conservations laws of physics i.e. con-

servation of baryon number which keeps all matter stable (just like kaon oscillation breaks strangeness quantum number). It also effectively makes neutron a Majorana fermion (albeit with a very tiny Majorana component), a point which is of some historical importance since in the original paper of Majorana, he contemplated neutron being the particle which is its own anti-particle.

The fact that $n - \bar{n}$ oscillation breaks baryon number requires us to first ascertain that the vast wealth of information available about the stability of matter from experiments and general cosmological observations allow for this process to have a strength that will make it observable with current technology. We show below that this is indeed the case.

Secondly as far as the physics implications go, another class of processes that probe baryon number nonconservation is proton decay[2, 3]. They have been the focus of experiment as well as theory for two reasons: first of all, simple grand unified theories[4] based on SU(5) and SO(10) predict its existence at a level not far from experimental capabilities; secondly, it used to be thought in the early 80's that these theories may also explain the origin of matter using proton decay as one of its key ingredients.

This situation however changed drastically with time due to two developements.

(a) The discovery of neutrino masses was confirmed in 1998 and a popular paradigm for the origin of their smallness became the seesaw mechanism[5], where new symmetry, B-L respected by then standard model forces had to be broken[1]. Note that the dominant proton decay modes of interest $p \to e^+\pi^o$ respect the $B - L$ symmetry and is therefore nor suited for the study of forces responsible for neutrino mases. On the other hand, $n - \bar{n}$ oscillations break B-L symmetry by two unjits exactly like the seesaw mechanism for neutrino masses does and therefore is uniquely suited for the study of the detailed nature as well as the

[1]Without any new symmetry, the seesaw scale would naturally be expected to be the Planck scale, which yields too small neutrino masses. A simple way to understand why the seesaw scale is lower than the Planck scale is to assume that it is associated with a symmetry e.g. B-L .

scale of seesaw mechanism. There is a large class of interesting gauge models where indeed neutrino Majorana masses via the seesaw mechanism leads directly to $n - \bar{n}$ oscillations.

(b) The second development is that a new mechanism for understanding the origin of matter was proposed in mid 80's that did not use proton decay but rather used the seesaw mechanism for neutrino masses[6] to generate matter-anti-matter asymmetry where again B-L symmetry played a key role. Although grand unified theories such as SO(10) could incorporate the B-L symmetry, these two developments took a lot of the wind from the argument in favor of proton decay being necessarily the most theoretically relevant mode for baryon nonconservation.

Furthermore, when the seesaw mechanism is embedded into GUT theories that include B-L symmetry, they generally tend to fix the scale of B-L symmetry breaking to be at the GUT scale, while there is no compelling reason for the seesaw scale to be that high. It could even be at the TeV scale without making the theory too unnatural. The seesaw scale could not therefore be probed by searching for proton decay whereas as already noted, in a large class of interesting theories, $n - \bar{n}$ oscillation does.

The question then arises as to whether all physics beyond the standard model is encapsulated in grand unified theories with neutrino masses being also a signal of the same GUT scale or whether the neutrino masses indicate an alternate path that contains rich physics far below the GUT scale. As we show below, $n - \bar{n}$ oscillation provides an effective probe of the second possibility which envisions a rich new layer of physics at intermediate scales that involves low B-L breaking. Theoretical models indicate that a wide range of scales, anywhere from a few TeV to $10^{11}$ GeV as well as the associated new physics could be probed by pushing the $n - \bar{n}$ oscillation oscillation time by three orders of magnitude over the current experimental lower limit. In particular, its discovery will rule out simple grand unified theories or in the very least have significant affect on their detailed nature. Furthermore

it will have profound effect on the nature of physics beyond the standard model as well as provide a new microphysics view for the origin of matter, different from leptogenesis. It also provides rich new physics that can be probed at the Large Hadron Collider.

The bottom line is therefore that it is urgent to pursue a dedicated search for for $n - \bar{n}$ oscillation that can push the sub-GUT physics frontier to a new level.

This review is organized into the following parts: in the first part, I discuss the phenomenon, its connection to nuclear instability ; in the next section, we discuss the broad character of microphysics that would lead to $n-\bar{n}$ oscillation and the kind of scales that can be probed by this. In the next section, the nature of physics beyond standard model probed by $n - \bar{n}$ oscillation will be discussed including a discussion as to why B-L is an important symmetry to be probed. In this section, we discuss specific and plausible gauge models for $n - \bar{n}$ oscillation both with and without supersymmetry. In the final section, we discuss the implication of observable $n - \bar{n}$ oscillation for baryogenesis and show how one can have a new class of models for late baryogenesis after electroweak phase transition and sphaleron freeze-out.

## 2 Phenomenology of $n - \bar{n}$ oscillation

In order to discuss the main features of $n - \bar{n}$ oscillation in various environments, it is sufficient to consider a simple $2 \times 2$ Hamiltonian for the evolution of an initial beam of slow moving neutrons and evaluate the probability of finding an anti-neutron in this beam after a time $t$.

$$i\hbar \frac{\partial}{\partial t} \begin{pmatrix} n \\ \bar{n} \end{pmatrix} = \begin{pmatrix} E_n & \delta m \\ \delta m & E_{\bar{n}} \end{pmatrix} \begin{pmatrix} n \\ \bar{n} \end{pmatrix} \tag{1}$$

where $\delta m$ denotes the $n-\bar{n}$ mixing which parameterizes at the nucleon level the underlying physics that breaks baryon number by two units. We will give examples of models where

this can happen at the appropriate level in a subsequent section. Note that we have left the energies of the $n$ and $\bar{n}$ arbitray since they could be affected in different ways in the presence of an external environment such as the nuclear field or a magnetic field etc.

The exact expression for the probability to find an antineutron at time given that at $t = 0$, $P_n(0) = 1$ and $P_{\bar{n}}(0) = 0$ is:

$$P_{\bar{n}}(t) \;=\; \frac{4\delta m^2}{\Delta E^2 + 4\delta m^2} sin^2(\sqrt{\Delta E^2 + 4\delta m^2})t; \tag{2}$$

where $\Delta E \;=\; E_n - E_{\bar{n}}$. The fact that neutron decays does not have any effect on this formula as long as the neutron flight time is much smaller than the life time of neutron.

Let us consider two extreme cases of this formula which also happen to be interesting for experimental purposes. For $\delta m \ll \Delta E$, this expression reduces to:

$$P_{\bar{n}} \sim \left(\frac{\delta m}{\Delta E_n}\right)^2 sin^2 \Delta E_n t \tag{3}$$

There are two special cases of this realizable in nature:

- Case (i): $\Delta E t \ll 1$: In this case we have, $P_{n\to\bar{n}} \sim (\delta m \cdot t)^2 \equiv \left(\frac{t}{\tau_{n-\bar{n}}}\right)^2$ This case corresponds to free neutron oscillation in vacuum.

- Case (ii): $\Delta E \cdot t \gg 1$: $P_{n\to\bar{n}} \sim \frac{1}{2}\left(\frac{\delta m}{2\Delta E_n}\right)^2$ This for instance corresponds to bound neutrons inside nucleus "oscillating" to anti-neutrons .

## 2.1 Stability of Nuclei and limit on $\tau_{n-\bar{n}}$:

As with all baryon number violating interactions, the existence of $n - \bar{n}$ oscillation will lead to nuclear instability. This will happen via a conversion of neutron to an anti-neutron inside the nucleus which will then annihilate with the surrounding nucleons and lead to instability. Clearly, if the nuclear transition probability inside the nucleus and in vacuum were equal,

this would cause all matter to disappear in a much shorter time than the age of the Universe and would be unacceptable. However, it is an experimental fact that inside the nucleus the neutron and the anti-neutron experience different nuclear potentials. In fact the difference is so large ($E_N - E_{\bar{N}} \sim 100$ MeV or so) that the transition probability formula relevant for this case is that in case (ii) above and leads indeed to nuclear instability lifetimes of order $10^{32}$ years or so. Careful and detailed analysis of this question has been done by Dover, Gal and Richards[11], Alberico et al[12] and more recently by Gal and collaborators[13]. Below we present a crude representation of the above works to get the basic physics across:

$$P^{nuc.}_{n-\bar{n}} \;\sim\; P_{n\to\bar{n}} \sim \left(\frac{\delta m}{2\Delta E_n}\right)^2. \tag{4}$$

where $P_{n\to\bar{n}}$ represents the free neutron oscillation probability in vacuum and $\delta m$ is the $\Delta B = 2$ off diagonal matrix element in the $n-\bar{n}$ mass matrix in Eq. (1) and is related to the oscillation time $\tau_{n-\bar{n}} = \frac{h}{2\pi\delta m}$. Present ILL limit on $\tau_{n-\bar{n}} \geq 10^8$ sec. translates to $\delta m \leq 6 \times 10^{-33}$ GeV. A crude estimate of the nuclear instability life time in the presence $n-\bar{n}$ oscillation from these intuitive considerations is $\tau_{Nucl.} \sim \left(\frac{\delta m}{2\Delta E_n}\right)^{-2} 10^{-23}$ sec.$\simeq 10^{34}$ years. More careful recent calculations that take into account the nuclear effects[13] are stated in terms of the following relation:

$$\tau_{Nuc.} \;=\; R(\tau_{n\bar{n}})^2 \tag{5}$$

where $R \sim 0.3 \times 10^{24}$ sec.$^{-1}$, which yields $\tau_{Nuc.} \sim 3 \times 10^{32}$ yrs. Present experimental limits on $\tau_{Nucl.}$ from various nucleon decay searches e.g. at Kamiokande, Sudan, IMB, Super-Kamiokande, SNO are given below in table I with the value of $\tau_{n-\bar{n}}$ noted there[16].

| Expt | source of neutrons | $\tau_{Nucl.}(yrs)$ | $\tau_{osc.}$(sec) |
|---|---|---|---|
| Soudan | $^{56}Fe$ | $0.72 \times 10^{32}$ | $1.3 \times 10^8$ |
| Frejus | $^{56}Fe$ | $0.65 \times 10^{32}$ | $1.2 \times 10^8$ |
| Kamiokande | $^{16}O$ | $0.43 \times 10^{32}$ | $1.2 \times 10^8$ |
| Super-K | $^{16}O$ | $1.77 \times 10^{32}$ | $2.3 \times 10^8$ |




Figure 1: Typical horizontal reactor set-up for $n - \bar{n}$ oscillation search

.

Table Caption: Experimental lower limits on $\tau_{n-\bar{n}}$ and the nuclei used for deducing the limit.

One can also anticipate a similar bound from the SNO experiment[20]. This implies $\rightarrow \delta m \leq 10^{-29}$ MeV. The nucleon decay searches for $\tau_{n-\bar{n}}$ become less efficient due to atmospheric neutrino induced background as we move to higher precision goals for $\tau_{n-\bar{n}}$. It would therefore appear that in order to extend the $\tau_{n-\bar{n}}$ search much beyond the current limit, one needs to focus more on free neutron oscillation experiments rather than than nucleon decay search.

We will see that present reactor neutron fluxes are precisely in the right range to probe these values of $\tau_{n-\bar{n}}$ that is compatibitible with matter stability as well as latest limits on proton decay.

## 2.2 Reactor Experiment to search for neutron oscillation

The basic idea behind a reactor search for $n-\bar{n}$ oscillation is to have a cold neutron beam from a reactor (neutron speed $v \sim 1-2$ kilo meters/sec.) and have the beam pass through a high vacuum, demagnetized region for as long as feasible to a detector at the end of this "tunnel". If one of the neutrons in the beam oscillates to an anti-neutron, this will annihilate in the detector and create a multi-pion signal with total energy of $\sim 2$ GeV. The need for demagnetization comes from the fact that the magnetic field of the Earth ($\sim 0.5$ Gauss) will split the neutron and anti-neutron energy levels by $\mu_n B \sim 3 \times 10^{-21}$ GeV. If we take a typical flight time in the "tunnel" of about one sec., it will imply $\mu_n B tsim 3 \times 10^{-21}$ GeV sec which about a 1000 times $\hbar$ and hence does not satisfy the condition for free neutron oscillation. It is therefore necessary to reduce the magnetic field to the level of $10^{-4}-10^{-5}$ Gauss. This is known technology that uses $\mu$-metal shielding. The typical horizontal set-up for such an experiment is given in Fig. 1 above. Under these conditions, one gets for the number of anti-neutrons for a given neutron beam the figure of merit of the experiment:

$$\#of events \;=\; \Phi_n \left(\frac{t}{\tau_{N-\bar{N}}}\right)^2 \times T \tag{6}$$

where $\Phi_n$ is the reactor flux; $v_n t$= distance to detector; $T$ running time. Maximum reactor fluxes for a 100 MW reactor is about $\sim 10^{13}-10^{14}$ neutrons per sec.; for $t = 0.1$ sec. and $T \sim 3$ years can yield a limit of $10^{10}$ sec. Using this technique, the first $n-\bar{n}$ search was carried out at the ILL laboratory in Grenoble, France and a lower limit of $\tau_{n-\bar{n}} \geq 8.6 \times 10^7$ sec. was obtained[15]. New searches have recently been proposed which could improve this limit by two orders of magnitude[17]. Fig. 2 compares the free oscillation time to the nuclear instability life time (figure courtesy of Y. Kamyshkov Ref.[17]). It is clear that time of flight $t$ anf the flux are two crucial factors that determine how precisely one can measure the $n-\bar{n}$ transition time. Clearly, the slower the neutrons the bigger is the time of flight. A possibility that is being pursued is to use ultracold neutrons with $\sim 100$ nano-eV kinetic




Figure 2: :
Comparision of free neutron oscillation time vs nuclear instability life time

energies which move with speeds of order 4-5 m/sec and are trapped in a bottle[18] and may become competitive with cold reactor beam experiments in near future[19].

# 3 Operator analysis of $\Delta B = 2$ processes and new physics scale probed

Whenever a new process is used to probe physics beyond the standard model, a general question that is essential to know is the mass scale probed by it. A simple way to get a general idea about the scale probed by a physical process is to do an operator analysis i.e. assuming a particular symmetry and spectrum of the low energy theory, write effective higher dimensional operators for the process under consideration and see for what value of the mass scaling the operator, the process is observable. This argument has been a useful tool to probe physics at short distance scales probed by processes such as proton decay, neutrino mass as well as corrections to standard model observables.

The method however has its limitations. For instance, often in these discussions, one uses the SM fermion spectrum but if there is a SM non-singlet new particle with a 100 GeV mass not discovered yet, the naive scale arguments can be misleading since new operators may appear. Similarly, the presence of unknown higher symmetries could also invalidate these arguments. We will apply this discussion to the $\Delta B = 2$ operators below to see the scale probed by neutron-anti-neutron oscillation.

## 3.1 Standard model operator

In the standard model, the effective operator responsible for $\Delta B = 2$ and $\Delta L = 0$ processes is given by:

$$\mathbf{O}_{\Delta B=2} = \frac{1}{M^5}[QQQQ(d^c d^c)^* + u^c d^c d^c u^c d^c d^c] \tag{7}$$

The strength of this operator $G_{\Delta B=2} \simeq \frac{1}{M^5}$. The $n-\bar{n}$ mixing mass can be deduced from this operator by simple dimensional analysis to be:

$$\delta m_{n-\bar{n}} \simeq c G_{\Delta B=2} \Lambda_{QCD}^6 = c \frac{\Lambda_{QCD}^6}{M^5} \tag{8}$$

where we chooze $c$ to be of order one. Attempts have been made in the past to estimate $c$ using bag as well as other phenomenological models for hadrons[21]. Taking the best current lower bound on $\tau_{n\leftrightarrow\bar{n}}$ from ILL reactor experiment [15] which is $10^8$ sec. and comparable bounds from nucleon decay search experiments [16], one can then obtain a lower limit on $M$ to be around 10-300 TeV depending on other couplings in a theory. If proposals to improve the precision of this search by at least two orders of magnitude [17] are carried out, , then in the context of the standard model particles, the mass scale probed will be around a 1000 TeV. Thus we see that neutron-anti-neutron oscillation will probe physics at scales far below the GUT scales that proton decay will probe. An important question to ask is whether the scale probed can be higher in the presence of new physics around the TeV scale. We discuss this below.

### 3.2 Supersymmetry and enhancement of $\Delta B = 2$ operator

In the presence of supersymmetry at the TeV scale, there are new particle such as squarks and sleptons with TeV scale mass. They can then enter the effective $\Delta B = 2$ operator. This kind of effect is familiar from discussions of supersymmetric grand unified theories where the presence of super-partners with TeV scale masses can drastically alter the operator analysis for nucleon decay. For instance the dimension six operator responsible for proton decay in non-susy models goes like $QQQL/M^2$ whereas the presence of squarks with TeV or sub-TeV masses, the dominant operator becomes $QL\tilde{Q}\tilde{Q}/M$, which has reduced power dependence on mass and is well known to put severe constraints on grand unified theories.

A typical leading operator for $n-\bar{n}$ oscillation for supersymmetric case is:

$$\mathbf{O}_{\Delta B=2} \;=\; \frac{1}{M^3} u^c d^c \tilde{d}^c \tilde{u}^c \tilde{d}^c \tilde{d}^c \tag{9}$$

Note that the power dependence on the seesaw scale (or B-L breaking scale) has now considerably softened. The conversion of susy particles to SM particles brings some suppression; but the the overall impact is that simple power counting arguments change. They imply that if this is the leading operator one can probe the seesaw scale upto $10^8$ GeV with $n-\bar{n}$ oscillation times of $10^{10}-10^{11}$ sec. The power dependence in fact softens even further if there are new color sextet particles at or below the TeV scale. For instance, if there is a scalar diquark coupled sextet field of $\Delta_{u^c u^c}$ type, the leading operator becomes:

$$\mathbf{O}_{\Delta B=2} \;=\; \frac{1}{M^2} d^c d^c \Delta_{u^c u^c} \tilde{d}^c \tilde{d}^c \tag{10}$$

and the scale reach of $n-\bar{n}$ goes up to $10^{11}$ GeV. If on the other hand there is a field $\Delta_{u^c d^c}$, the leading order operator becomes:

$$\mathbf{O}_{\Delta B=2} \;=\; \frac{1}{M} d^c d^c \Delta_{u^c d^c} \Delta_{u^c d^c} \tag{11}$$

increasing the scale reach to the GUT scale. The question then arises whether there are plausible models where this can happen. Below I give an example of models which have TeV scale $\Delta_{u^c u^c}$ fields naturally so that an $1/M^2$ dependence on the seesaw scale is quite natural. From this, we see that scale of observable $n-\bar{n}$ physics can be anywhere from 300 TeV to $10^13$ TeV

# 4 Neutrino mass physics probed by $n-\bar{n}$ oscillation

To see what new physics is revealed by the search for $n-\bar{n}$ oscillation, we give several examples where low scale for seesaw manifests in the non-leptonic sector of the theory as

$n - \bar{n}$ oscillation. This happens due to the connection of seesaw to local or global B-L symmetry. There could also be other kind of beyond the standard model scenarioss where extra space dimensions curled up to $\text{TeV}^{-1}$ length scales and where additional dynamical inputs can lead to observable $n - \bar{n}$ oscillation. In the former case, it will throw important light on the physics associated with neutrino mass and we discuss this below.

There are various reasons to think that B-L symmetry is an intimate feature of physics beyond the standard model. Some of them are as follows:

- (i) The seesaw mechanism for understanding small neutrino masses requires the introduction of right handed neutrinos[5]. In the presence of three right handed neutrinos it is more natural for B-L symmetry which was a global symmetry of the SM Lagrangian to become a local symmetry;

- (ii) An inherent aspect of seesaw is the Majorana mass of the right handed neutrino that breaks the B-L symmetry providing one way to understand why the seesaw scale is so much less than the Planck scale;

- (iii) A third reason appears once one admits the presence of supersymmetry at the TeV scale, as is widely believed and requires the lightest SUSY particle (LSP) to be naturally stable in order to be the dark matter candidate of the universe. The simplest way to have SUSY LSP naturally stable is to have B-L as a symmetry of physics beyond MSSM[7].

If indeed B-L symmetry is present in nature, there are several questions that immediately come to mind:

- (a) Is it a global or local symmetry ?

- (b) is it a broken or unbroken symmetry ?

- (c) if it is broken, what is the breaking scale and what is the associated physics ?

I will now argue that search for the process of neutron-anti-neutron oscillation[8, 9, 10] can provide a partial answers to some of these important questions: in particular the question of scale of B-L breaking and associated expanded gauge symmetry and hence physics associated with the origin of neutrino mass. The point is that since a $\Delta B = 2$ operator breaks B-L by two units, it is natural to associate the scale associated with the $\Delta B = 2$ operator also with the scale of B-L breaking since neither $B$ nor $L$ are separate symmetries of the standard model but rather the symmetry is $B-L$ . Since seesaw mechanism also breaks B-L symmetry, $M$ could also be the seesaw scale. In fact, in the context of models such as those based on $SU(2)_L \times SU(2)_R \times SU(4)_c$[14], the same mass scale $M$ is responsible for both processes as was first shown in Ref.[10]. Another purely group theoretical way to see this is to note that in left-right symmetric models, there is a relation between the electric charge and the B-L generator as follows:

$$Q \;=\; I_{3L} \;+\; I_{3R} \;+ \frac{B-L}{2} \tag{12}$$

For distance scales shorter than the electroweak scale, we have $\Delta I_{3L} = 0$ and electric charge conservation then implies that $\Delta I_{3R} \;=\; -\frac{B-L}{2}$ implying that parity violation (or $\Delta I_{3R} \;=\; 1$) implies not only that $\Delta L \;=\; 2$ i.e. seesaw mechanism for neutrino mass but in the non-leptonic sector it implies $\Delta B \;=\; 2$ which is of course $n - \bar{n}$ oscillation.

Thus the search for $n-\bar{n}$ oscillation could not only illuminate the nature of the important symmetry, B-L but could also be one way to unravel the mystery of the seesaw mechanism that is expected to be major player in the physics of neutrino mass as well as the origin of parity violation in Nature.

# 5 $SU(2)_L \times SU(2)_R \times SU(4)_c$ model with light diquarks

In this section we demonstrate by an explicit example how neutrino mass and $n - \bar{n}$ are intimately connected in this partial unification theory. We first recapitulate some elementary facts about the model.

The quarks and leptons in this model are unified and transform as $\psi : (\mathbf{2}, \mathbf{1}, \mathbf{4}) \oplus \psi^c : (\mathbf{1}, \mathbf{2}, \bar{\mathbf{4}})$ representations of $SU(2)_L \times SU(2)_R \times SU(4)_c$. For the Higgs sector, we choose, $\phi_1 : (\mathbf{2}, \mathbf{2}, \mathbf{1})$ and $\phi_{15} : (\mathbf{2}, \mathbf{2}, \mathbf{15})$ to give mass to the fermions. The $\Delta^c : (\mathbf{1}, \mathbf{3}, \mathbf{10}) \oplus \bar{\Delta}^c : (\mathbf{1}, \mathbf{3}, \overline{\mathbf{10}})$ to break the $B - L$ symmetry. There are color sextet scalar diquark fields contained in the $\Delta^c : (\mathbf{1}, \mathbf{3}, \mathbf{10})$ multiplet, which will play a crucial role in generating $n - \bar{n}$ oscillation. We first discuss the non-0supersymmetric version of the model in next subsection and follow it up by the supersymmetric generalization where the scale reach of $n - \bar{n}$ goes up all the way to $10^{10}$ GeV or so.

## 5.1 Non-supersymmetric version

The Higgs potential for this model can be written as :

$$V(\phi_0, \phi_{15}, \Delta_{R,L}) \;=\; V_0 \;+\; V_m \tag{13}$$

where

$$V_m \;=\; \lambda_m \epsilon^{\mu\nu\sigma\rho} \epsilon^{\mu'\nu'\sigma'\rho'} \Delta_{R,\mu\mu'} \Delta_{\nu\nu'} \Delta_{\sigma\sigma'} \Delta_{\rho\rho'} + \; (R \leftrightarrow L) + \; h.c. \tag{14}$$

This term is crucial for generating baryon number non-conservation in this theory and without it, the model cannot explain the origin of matter in the universe (as we see below). We will see that the same term is responsible for the $\Delta B = 2$ $n - \bar{n}$ oscillation.

In order to discuss neutrino masses and $n - \bar{n}$ oscillation, let us write down the Yukawa

couplings in the model:

$$L_Y \;=\; h_1\bar{\Psi}_L\Phi_1\Psi_R \;+\; h_{15}\bar{\Psi}_L\Phi_{15}\Psi_R \;+\; f\Psi_R^T C^{-1}\Delta_R\Psi_R \;+(R\leftrightarrow L)+H.C. \tag{15}$$

Using standard spontaneous breaking via the vev of $(1,3,10)$ to the standard model gauge group and $\Phi_{1,15}$ to give mass to both to charged fermions and Dirac mass to neutrinos, we implement the seesaw mechanism for neutrinos. It was shown in ref.[10] that the same $(1,3,10)$ via the potential term $V_m$ leads to $n-\bar{n}$ oscillation via the diagram in Fig. 3. The




Figure 3: A Feynman diagram contributing to $n-\bar{n}$ oscillation as discussed in Ref. 8. Another class of graphs involves two $\Delta_{u^cd^c}$ and one $\Delta_{d^cd^c}$ scalar exchanges.

strength of $n-\bar{n}$ oscillation is given by:

$$G_{\Delta B=2} \;=\; \frac{\lambda f_{11}^3 v_{BL}}{M^6_{\Delta_{qq}}} \tag{16}$$

We see the fifth power dependence on the seesaw scale since the diquarks are expected to be

at seesaw scale by the usual naturalness considerations. Also note that even after symmetry breaking this model has a $Z_2$ symmetry given by $(-1)^B$.

It has recently been shown[22] that this model also explains the baryon asymmetry of the universe via the mechanism of post-sphaleron baryogenesis to be discussed below[23].

## 5.2 Supersymmetric version

We now discuss the supersymmetric version of this model where the triplet fields doubled to cancel the gauge anomalies. We also add a $B-L$ neutral triplet $\Omega : (\mathbf{1}, \mathbf{3}, \mathbf{1})$ which helps to reduce the accidental global symmetry of the model and hence the number of light diquark states. The superpotential of this model is given by:

$$W \;=\; W_Y \;+\; W_{H1} + \;W_{H2} + \;W_{H3} \tag{17}$$

where

$$W_{H1} \;=\; \lambda_1 S(\Delta^c\bar{\Delta}^c - M^2) \;+ \lambda_C \Delta^c\bar{\Delta}^c\Omega \;+ \mu_i \mathrm{Tr}\,(\phi_i\phi_i) \tag{18}$$

$$W_{H2} \;=\; \lambda_A \frac{(\Delta^c\bar{\Delta}^c)^2}{M_{P\ell}} + \lambda_B \frac{(\Delta^c\Delta^c)(\bar{\Delta}^c\bar{\Delta}^c)}{M_{P\ell}} \tag{19}$$

$$W_{H3} \;=\; \lambda_D \frac{\mathrm{Tr}\,(\phi_1\Delta^c\bar{\Delta}^c\phi_{15})}{M_{P\ell}}\,, \tag{20}$$

$$\tag{21}$$

$$W_Y \;=\; h_1\psi\phi_1\psi^c + h_{15}\psi\phi_{15}\psi^c + f\psi^c\Delta^c\psi^c. \tag{22}$$

Note that since we do not have parity symmetry in the model, the Yukawa couplings $h_1$ and $h_{15}$ are not symmetric matrices. When $\lambda_B = 0$, this superpotential has an accidental global symmetry much larger than the gauge group[25]; as a result, vacuum breaking of the $B-L$

symmetry leads to the existence of light diquark states that mediate $N \leftrightarrow \bar{N}$ oscillation and enhance the amplitude. In fact it was shown that for $\langle \Delta^c \rangle \sim \langle \bar{\Delta}^c \rangle \neq 0$ and $\langle \Omega \rangle \neq 0$ and all VEVs in the range of $10^{11} - 10^{12}$ GeV, the light states are those with quantum numbers: $\Delta_{u^c u^c}$. The symmetry argument behind is that [25] for $\lambda_B = 0$, the above superpotential is invariant under $U(10,c) \times SU(2,c)$ symmetry which breaks down to $U(9,c) \times U(1)$ when $\langle \Delta^c_{\nu^c \nu^c} \rangle = v_{BL} \neq 0$. This results in 21 complex massless states; on the other hand these vevs also breaks the gauge symmetry down from $SU(2)_R \times SU(4)_c$ to $SU(3)_c \times U(1)_Y$. This allows nine of the above states to pick up masses of order $g v_{BL}$ leaving 12 massless complex states which are the six $\Delta^c_{u^c u^c}$ plus six $\bar{\Delta}^c_{u^c u^c}$ states. Once $\lambda_B \neq 0$ and is of order $10^{-2} - 10^{-3}$, they pick up mass (call $M_{u^c u^c}$) of order of the elctroweak scale.

### 5.3 A new diagram for neutron-anti-neutron oscillation

To discuss $n \leftrightarrow \bar{n}$ oscillation, we introduce a new term in the superpotential of the form[10]:

$$W_{\Delta B=2} \;=\; \frac{1}{M_*} \epsilon^{\mu' \nu' \lambda' \sigma'} \epsilon^{\mu \nu \lambda \sigma} \Delta^c_{\mu \mu'} \Delta^c_{\nu \nu'} \Delta^c_{\lambda \lambda'} \Delta^c_{\sigma \sigma'} \,, \tag{23}$$

where the $\mu, \nu$ etc stand for $SU(4)_c$ indices and we have suppressed the $SU(2)_R$ indices. Apriori $M_*$ could be of order $M_{P\ell}$; however the terms in Eq.(2) are different from those in Eq. (4); so they could arise from different a high scale theory. The mass $M_*$ is therefore a free parameter that we choose to be much less than the $M_{P\ell}$. This term does not affect the masses of the Higgs fields. When $\Delta^c_{\nu^c \nu^c}$ acquires a VEV, $\Delta B = 2$ interaction are induced from this superpotential, and $n \leftrightarrow \bar{n}$ oscillation are generated by two diagrams given in Fig. 3 and 4. The first diagram (Fig. 3) in which only diquark Higgs fields are involved was already discussed in [10] and goes like $G_{n \leftrightarrow \bar{n}} \simeq \frac{f_{11}^3 v_{BL} M_\Delta}{M^2_{u^c u^c} M^4_{d^c d^c} M_*}$, Taking $M_{u^c u^c} \sim 350$ GeV, $M_{d^c d^c} \sim \lambda' v_{BL}$ and $M_\Delta \sim v_{BL}$ as in the argument [25], we see that this diagram scales like $v_{BL}^{-3} v_{wk}^{-2}$. In ref.[24] a new diagram (Fig. 4) was pointed out which owes its origin to

supersymmetry. We get for its contribution to $G_{\Delta B=2}$:

$$G_{n\leftrightarrow\bar{n}} \simeq \frac{g_3^2}{16\pi^2}\frac{f_{11}^3 v_{BL}}{M^2_{u^c u^c} M^2_{d^c d^c} M_{\rm SUSY} M_*}. \tag{24}$$

Using the same arguments as above, we find that this diagram scales like $v_{BL}^{-2} v_{wk}^{-3}$ which is therefore a significant enhancement over diagram in Fig.3 when $v_{BL} \gg v_{wk}$. In order to




Figure 4: The new Feynman diagram for $n - \bar{n}$ oscillation.

estimate the rate for $n \leftrightarrow \bar{n}$ oscillation, we need not only the different mass values for which we now have an order of magnitude, we also need the Yukawa coupling $f_{11}$. Now $f_{11}$ is a small number since its value is associated with the lightest right-handed neutrino mass.

However, in the calculation we need its value in the basis where quark masses are diagonal. We note that the $n-\bar{n}$ diagrams involve only the right-handed quarks, the rotation matrix need not be the CKM matrix. The right-handed rotations need to be large e.g. in order to involve $f_{33}$ (which is $O(1)$), we need $(V_R^{(u,d)})_{31}$ to be large, where $V_L^{(u,d)\dagger} Y_{u,d} V_R^{(u,d)} = Y_{u,d}^{\rm diag.}$. The left-handed rotation matrices $V_L^{(u,d)}$ contribute to the CKM matrix, but right-handed rotation matrices $V_R^{(u,d)}$ are unphysical in the standard model. In this model, however, we get to see its contribution since we have a left-right gauge symmetry. Let us now estimate the time of oscillation. When we start with a $f$-diagonal basis (call the diagonal matrix $\hat{f}$), the Majorana coupling $f_{11}$ in the diagonal basis of up- and down-type quark matrices can be written as $(V_R^T \hat{f} V_R)_{11} \sim (V_{31}^R)^2 \hat{f}_{33}$. Now $\hat{f}_{33}$ is $O(1)$ and $V_{31}^R$ can be $\sim 0.6$, so $f_{11}$ is about 0.4 in the diagonal basis of the quark matrices. We use $M_{\rm SUSY}$, $M_{u^c u^c} \sim 350$ GeV and $v_{BL} \sim 10^{12}$ GeV. The mass of $\tilde{\Delta}_{d^c d^c}$ i.e. $M_{d^c d^c}$ is $10^9$ GeV which is obtained from the VEV of $\Omega : (\mathbf{1}, \mathbf{3}, \mathbf{1})$. We choose $M_* \sim 10^{13}$ GeV. Putting all the above the numbers together, we get

$$G_{n \leftrightarrow \bar{n}} \simeq 1 \cdot 10^{-30}\ {\rm GeV}^{-5}. \tag{25}$$

Taking into account the hadronic matrix element effect, the $n-\bar{n}$ oscillation time is found to be about $2.5 \times 10^{10}$ sec which is within the reach of possible next generation measurements. If we chose, $M_* \simeq M_{P\ell}$, we will get for $\tau_{n-\bar{n}} \sim 10^{15}$ sec. unless we choose the $M_{d^d d^c}$ to be lower (say $10^7$ GeV). This is a considerable enhancement over the nonsupersymmetric model of [10] with seesaw scale of $10^{12}$ GeV. We also note that as noted in [10] the model is invariant under the hidden discrete symmetry under which a field $X \to e^{i\pi B_X} X$, where $B_X$ is the baryon number of the field $X$. As a result, proton is absolutely stable in the model. Furthermore, since R-parity is an automatic symmetry of MSSM, this model has a naturally stable dark matter.

# 6 Baryogenesis and $n - \bar{n}$ oscillation

In the early 1980's when the idea of neutron-anti-neutron oscillation was first proposed in the context of unified gauge theories, it was thought that the high dimensionality of the $\Delta B \neq 0$ operator would pose a major difficulty in understanding the origin of matter in the Universe. The main reason for this assessment is that the higher dimensional operators remain in thermal equilibrium until late in the evolution of the universe without contradicting any low energy observations. This is because the thermal decoupling temperature $T_*$ for such interactions goes roughly like $v_{BL}\left(\frac{v_{BL}}{M_P}\right)^{1/9}$ which can be in the range of temperatures where B+L violating sphaleron transitions are in full thermal equilibrium, say for example $v_{BL} \simeq 10 - 100$ TeV, as required for the case of observable $n - \bar{n}$ oscillation. They will therefore erase any baryon asymmetry generated in the very early moments of the universe (say close to the GUT time of $10^{-36}$ sec. or so) in then prevalent baryogenesis models. Even though GUT baryogenesis models are no more popular, the same argument will apply to any other high temperature baryogenesis mechanism.

In models with observable $n - \bar{n}$ oscillation therefore, one has to search for new mechanisms for generating baryons below the weak scale. In this section, we discuss such a possibility[23] which was discussed in a recent paper. As we see below, it is ideally suited for embedding into the $G_{224}$ model discussed in Ref.[10] and leads to an interesting side "bonus" that it puts an upper limit on the neutron-anti-neutron oscillation time $\tau_{n-\bar{n}}$.

As an illustration of the way the new mechanism operates, let us assume that there is a complex scalar field that couples to the $\Delta B = 2$ operator as follows i.e. $L_I = S u^c d^c d^c u^c d^c d^c / M^6$, where the scalar boson has mass of order of the weak scale and $B = 2$. When $< S > \neq 0$, this interaction leads to baryon number violation by two units and observable $n - \bar{n}$ transition if $M$ is in the few hundred to 1000 GeV range. Writing $S = \frac{1}{\sqrt{2}}(v_{BL} + S_r + S_I)$, we see

that the direct decay of $S_r$ involves both $\Delta B = \pm 2$ final states and thereby satisfies the first requirement for baryogenesis.

The first point to note is that the high dimension of $L_I$ allows the scalar $\Delta B \neq 0$ decay to go out of equilibrium at weak scale temperatures. This satisfies the out of equilibrium condition given by Sakharov for generating matter-anti-matter asymmetry. To see this, let us give some examples: if we chose the proton decay operators such as $QQQL$, the decoupling temperature consistent with present experimental bounds on proton life time would be around $10^{15}$ GeV or so. So to apply our mechanism, we need to consider higher dimensional operators. A typical example of such an operator that we will focus on in this paper is of type: $u^c d^c d^c u^c d^c d^c / M^5$. As note earlier, this operator leads to $n - \bar{n}$ oscillation and for this to be in the observable range, the associated mass scale has to be in the few TeV range. The decupling temperature can then be easily estimated from the formula $T_d \sim M \left(\frac{10M}{M_P}\right)^{1/9} \sim 10^{-1.5} M$. For appropriate values of $M$ in the range of interest $T_d$ can be below 100 GeV so that the sphalerons have gone out of equilibrium and baryogenesis follows. To make these ideas more concrete, below we give an explicit example[23].

We consider an effective sub-TeV scale model that gives rise to the above higher dimensional operator for $n \leftrightarrow \bar{n}$ oscillation. It consists of the following color sextets, $SU(2)_L$ singlet scalar bosons $(X, Y, Z)$ with hypercharge $-\frac{4}{3}, +\frac{8}{3}, +\frac{2}{3}$ respectively that couple to quarks. These states emerge naturally if this low scale theory is embedded into the $SU(2)_L \times SU(2)_R \times SU(4)_c$ (denoted by $G_{224}$) model described in the previous section. The scalar field $S$ can be identified with the complex field $\Delta_{\nu^c \nu^c}$ whose vev breaks B-L symmetry by two units as is the case for us and $S_r \equiv Re(\Delta^0_{\nu^c \nu^c}$[22]. The $X, Y, Z$ fields can be identified with the color sextet partners of $\Delta_{\nu^c \nu^c}$ in the $(1, 3, 10)$ multiplet i.e. the $\Delta_{d^c d^c}, \Delta_{u^c u^c}$ and $\Delta_{u^c d^c}$ respectively. In the $X, Y, Z$ notation, one can write down the following standard model invariant interaction

Lagrangian:

$$L_I = h_{ij} X d_i^c d_j^c + f_{ij} Y u_i^c u_j^c + g_{ij} Z (u_i^c d_j^c + u_j^c d_j^c) + \lambda_1 S X^2 Y + \lambda_2 S X Z^2 \quad (26)$$

The complex scalar field $S$ clearly has $B - L = 2$. To see the constraints on the parameters of the theory, we note that the present limits on $\tau_{n-\bar{n}} \geq 10^8$ sec. implies that the strength $G_{n-\bar{n}}$ of the the $\Delta B = 2$ transition is $\leq 10^{-28}$ GeV$^{-5}$. From the above figure, we conclude that

$$G_{n-\bar{n}} \simeq \frac{\lambda_1 M_1 h_{11}^2 f_{11}}{M_Y^2 M_X^4} + \frac{\lambda_2 M_1 h_{11} g_{11}^2}{M_X^2 M_Z^2} \leq 10^{-28} GeV^{-5}. \quad (27)$$

For $\lambda_{1,2} \sim h_{11} \sim f_{11} \sim g_{11} \sim 10^{-3}$, we require $M_1 \sim M_{X,Y,Z} \simeq 1$ TeV to satisfy this experimental bound. In our discussion of generic models of this type, we will stay close to this range of parameters and see how one can understand the baryon asymmetry of the universe. The singlet field will play a key role in the generation of baryon asymmetry. We assume that $< S > \gg M_X$ and $M_{S_r} \sim 100 - 500$ GeV. It can then decay into final states with $B = \pm 2$ i.e. six quarks and six anti-quarks.

Note that the Lagrangian of Eq. (26), leads to tree level contribution to flavor changing neutral current processes such as $K - \bar{K}$, $B - \bar{B}$ mixings etc and that will restrict the form of the flavor structure of the coupling matrices suitably. These constraints can also be satisfied in a realistic $G_{224}$ embedded model which also explain the neutrino masses and mixing[22]. This is an important consideration since the magnitude of the baryon asymmetry depends on this detailed flavor structure.

On the way to calculating the baryon asymmetry, let us first discuss the out of equilibrium condition. As the temperature of the universe falls below the masses of the $X, Y, Z$ particles, the annihilation processes $X\bar{X} \to d^c \bar{d^c}$ (and analogous processes for $Y$ and $Z$) remain in equilibrium. As a result, the number density of $X, Y, Z$ particles gets depleted as $e^{-M_X/T}$

and only the $S_r$ particle survives along with the usual standard model particles. One of the primary generic decay modes of $S_r$ is $S_r \to u^c d^c d^c u^c d^c d^c$. There could be other decay modes which depend on the details of the model and one has to ensure that the rates to other modes are smaller than the six quark mode.

The generic chain of events leading to baryogenesis in this model is the following. At $T \sim M_{S_r}$, the decay rate is smaller than the Hubble expansion rate. As the Universe cools below this temperature, the decay rate remains constant whereas the expansion rate of the universe is slowing down. So at a temperature $T_d$ far below $M_S$, $S$ will start to decay when the decay rate $\Gamma \sim H$ with $T_d$ is given by:

$$T_d \simeq (\frac{M_{P\ell} M_S^{13}}{(2\pi)^9 M_X^{12}})^{1/2} \tag{28}$$

Since the corresponding epoch must be above that the QCD phase transition temperature, this puts a constraint on the parameters of the model. Typically, we need to have $M_S/M_X \sim 0.5$ or so due to high power dependence of $T_d$ on this ratio. This implies that the $X, Y, Z$ masses cannot be arbitrarily high, since the heavier these particles are, the lower $T_d$ will be. We expect this upper limit to be in the TeV range at most as we show below .

It is well known that baryon asymmetry can arise only via the interference of a tree diagram with a one loop diagram. The tree diagram is clearly the one where $S \to 6q$. There are however two classes of loop diagrams that can contribute: one where the loop involves the same fields $X, Y$ and $Z$. A second one involves W-exchange, which involves only standard model physics at this scale (Fig. 6). We find that the second contribution can actually dominate. In fact, in the $G_{224}$ embedding of the model, the first diagram vanishes. The second diagram also has the advantage that it involves less number of arbitrary parameters and the source of CP violation in this case is the same as the CKM CP violation present in the standard model. The observed baryon asymmetry is related to the primordial baryon

asymmetry $\epsilon_B$ as $\eta_B \simeq (\frac{g^*_{rec}}{g^*_{T_d}})^{3/4}\epsilon_B)$ where

$$\epsilon_B \simeq \frac{n_S}{n_\gamma}\frac{\Gamma(S \to 6q) - \Gamma(S \to 6\bar{q})}{\Gamma(S)} \tag{29}$$

We find that




Figure 5: Tree level Feynman diagram for S decays.




Figure 6: One loop diagram for S decays.

$$\frac{\epsilon_B^{\text{vertex}}}{\text{Br}} \simeq -\frac{\alpha_2}{4}\frac{\text{Im}\text{Tr}[gg^T\hat{M}_u V M_d g^\dagger M_u V^* \hat{M}_d]}{\text{Tr}(g^\dagger g)M_W^2 M_S^2} \ . \tag{30}$$

The magnitude of the asymmetry depends on $M_S$ as well as the detailed profile of the various coupling matrices $h, g, f$ and we can easily get the desired value of the baryon asymmetry by appropriately choosing them. An important consequence of the above equation regardless of the details is that if $M_S$ is much bigger than about 500 GeV, baryon asymmetry becomes

very small. This implies that $M_{X,Y,Z}$ in turn should not be much larger than a TeV, implying that they can be accessible at the LHC. This also implies that the strength of $n-\bar{n}$ transition has a lower limit or an upper limit on $\tau_{n-\bar{n}}$.

There is a dilution of the baryon asymmetry arising from the fact that $T_d \ll M_S$ since the decay of $S$ also releases entropy into the universe. Thius dilution factor is

$$d_B \simeq \epsilon_B \frac{(0.32 g_*(T_d) T_d)^{3/4}}{(0.12 M_S + 0.32 G_* T_D)^{3/4}} \tag{31}$$

Since the decay rate of the $S$ boson depends inversely as a high power of $M_{X,Y}$, higher $X, Y$ bosons would imply that $\Gamma_S \sim H$ is satisfied at a lower temperature and hence give a lower $d_B$. For our preferred choice of parameters i.e. $M_X \sim 2M_S \sim$ TeV, we find $d_B \sim 20\%$. Also for choice of the coupling parameters $\lambda \sim f \sim h \sim g \sim 10^{-3}$, and $M_S \simeq 200\ GeV$ we find $\tau_{n-\bar{n}} \leq 10^{10}\ sec.$

We also note that if the model is embedded into a $G_{224}$ group, all three coupling matrices $h, g, f$ become equal; there are also some constraints on the gauge boson spectrum in this model[22]. As noted above, an interesting consequence of adequate baryogenesis is that there are new sub-TeV- TeV scale color sextet scalar bosons which can be observed at LHC[26].

# 7 Extra dimensional models with observable $n - \bar{n}$ oscillation

The possibility that there may be extra compact space dimensions has been under extensive theoretical as well as experimental investigation inspired by the fact that superstring theories do predict the existence of such models. In a large class of interesting models, these dimensions are believed to be of order TeV$^{-1}$ size[27, 28] making their manifestations in many physical situations of experimental interest not only for colliders but also for low energy experiments. One class of such possibilities includes the breakdown of baryon number

by two units. We note two such discussions:

(i) In ref.[29], it was noted that if there are millimeter size extra dimensions, one way to understand the observed fermion mass hierarchies would be to distribute the standard model fermions in the bulk. In such a scenario, their Yukawa couplings would be suppressed if they are further apart[30]. Thus in the context of a 5- or 6 dimensional scenario, observed mass hierarchies pretty much fix the locations of the various standard model fermions in the bulk. One can then ask the question as to what happens to higher dimensional operators such as those giving proton decay and $n-\bar{n}$ oscillation. Since these are low scale models, one has to separate the quarks and leptons sufficiently to suppress these operators since their strength is given by the overlap integral of the various wave functions in the extra dimension. The $n-\bar{n}$ operator however involves only the right handed quarks and their locations is generally fixed by the mass hierarchy considerations. So there is no freedom to suppress their strengths. The strength of such operators was calculated in the context of six dimensional models in Ref.[29] and found to be in the observable range for extra dimension sizes of order $45-100$ TeV, which are of the same order as being contemplated for phenomenological viable models of this kind. These considerations could also be applied to the case Randall-Sundrum models where the SM fermions are also generally expected to be distributed in the bulk.

(ii) A second class of models were discussed in ref.[31], where general considerations of global charge separation from the branes is used to argue that one cannot have proton decay but observable $n-\bar{n}$ oscillations due to the fact that neutrons are both color and electric charge neutral. Such baryon number nonconservation occurs due to brane fluctuations which can create baby branes that carry baryon number if they are color and charge neutral.




Figure 7: neutron oscillation induced in R-parity breaking MSSM

# 8 R-parity breaking models

Finally, there have been explorations of $n - \bar{n}$ oscilation to supersymmetry. As is well known, in minimal supersymmetric extensions of the standard model (MSSM), a baryon number violating operator of the form $\lambda^{''} u^c d^c d^c$ is allowed in the superpotential. By itself it can lead to $\Delta B = 2$ operator of the form $(u^c d^c d^c)^2$ at tree level via gluino exchange (Figure 7). Typical strength of such operators arising from this new interaction is given[32] by:

$$G_{\Delta B=2} = \frac{4\pi\alpha_s \lambda^{''2}_{1\tilde{1}2}}{M^4_{\tilde{q}} M_{\tilde{G}}} \tag{32}$$

Note that due to color anti-symmetry, at the tree level, this interaction leads $\Lambda - \bar{\Lambda}$ oscillation rather than $n - \bar{n}$ oscillation. However once radiative corrections are included, this operator via flavor changing effects gives rise to neutron oscillation[33]. This puts constraints on the R-parity violating couplings of the $\lambda^{''}$ type. It must be noted that if these couplings are present in combination with R-P violating operators of type $QLd^c$, then it leads to catastrophic proton decay rates[34]; so in discussing $\Delta B = 2$ transitions, we are assuming that $QLd^c$ operators are absent.

# 9 Conclusion

In summary, there is now a widely held belief that neutrino mass provides a clue to the existence of a new local B-L symmetry of nature. A key question in particle theory beyond the standard model is then to find out the scale and detailed nature of physics associated

with this new symmetry. The situation is similar to the late 60's when the hints in favor of the existence of a fundamental local $SU(2)$ symmetry of weak interaction were becoming compelling that eventually led to the triumphant gauge revolution in the 1970's. They were subsequently confirmed by the discovery of neutral currents, followed by the discovery of the W and Z boson. The two paths before us right now are (i) whether the B-L forces responsible for neutrino mass represent physics close to the gravity scale of $10^{18}$ GeV as exemplified, say by the grand unified theories or (ii) scales multi-TeV range far below the GUT scale with signals visible to the Large Hadron Colliders. One direct way to explore this new physics is to to carry out an experimental search for $n-\bar{n}$ oscillation. As noted the discovery of $n-\bar{n}$ oscillation will have far reaching implications not only for the nature of forces and matter but will completely alter our thinking about such cosmological issues as the origin of matter, nature of dark matter etc.

The main points stressed in this review are that: (a)there are a wide class of models e.g. $SU(2)_L \times SU(2)_R \times SU(4)_c$ with type II seesaw, models based on TeV scale extra dimension models where $n-\bar{n}$ oscillation is in a range accessible to currently planned reactor experiments and they will extend the existing limit on the strength of this process to an exciting range; (b) Models that predict observable $n-\bar{n}$ oscillation provide a new way to understand the origin of matter where matter creation moment is "much" later than currently contemplated models such as leptogenesis or GUT scale baryogenesis. More importantly, unlike the other mechanisms for baryogenesis, this mechanism can be tested in current collider experiments as well as neutrino experiments in some specific realizations.

On the experimental front, searches for $n-\bar{n}$ transition using the proton decay type set-ups are back ground limited[35] and cannot be used to push the limits for $\tau_{n-\bar{n}}$ too much beyond the present limit. Also an interesting class of theories that gives rise to $n-\bar{n}$ induced baryogenesis lead to new coloe sextet particles observable at LHC[26].

Finally, it is also worth noting that there is a related phenomenon involving neutrons i.e. neutron-mirror neutron oscillation that can also be searched for in experimental set-ups similar to that searching for $n - \bar{n}$ oscillation[36]. This new class of oscillations have a much weaker experimental limit on its strength and could be improved by longer "baseline" searches being contemplated for $n - \bar{n}$ oscillation.

I would like to thank K. S. Babu for many discussions on theoretical aspects of the topic and Y. Kamyshkov, G. Greene, Mike Snow and Albert Young for many discussions on the experimental prospects for $n-\bar{n}$ oscillation. I am grateful to K. S. Babu and Y. Kamyshkov for carefully reading the manuscript and suggesting improvements. This work is supported by the National Science Foundation grant no. Phy-0652363.

# References

[1] For a review, see R. N. Mohapatra, Nucl. Instrum. Meth. A **284**, 1 (1989).

[2] For recent reviews and references, see P. Nath and P. Fileviez Perez, Phys. Rept. **441**, 191 (2007) and Kaladi S. Babu, Talk at the BNL workshop on "Underground Detectors Investigating Grand Unification" http://www.bnl.gov/udig/.

[3] For discussion of both proton decay and neutron oscillation search possibilities at DUSEL, see S. Raby *et al.*, "DUSEL Theory White Paper," arXiv:0810.4551 [hep-ph].

[4] For review of grand unified theories, see S. Raby, in *The Review of Particle Physics* C. Amsler et al., Physics Letters B667, 1 (2008).

[5] P. Minkowski, Phys. lett. B **67**, 421 (1977); M. Gell-Mann, P. Ramond, and R. Slansky, *Supergravity* (P. van Nieuwenhuizen et al. eds.), North Holland, Amsterdam, 1979, p. 315; T. Yanagida, in *Proceedings of the Workshop on the Unified Theory and the*

*Baryon Number in the Universe* (O. Sawada and A. Sugamoto, eds.), KEK, Tsukuba, Japan, 1979, p. 95; S. L. Glashow, *The future of elementary particle physics*, in *Proceedings of the 1979 Cargèse Summer Institute on Quarks and Leptons* (M. Lévy et al. eds.), Plenum Press, New York, 1980, p. 687; R. N. Mohapatra and G. Senjanović, Phys. Rev. Lett. **44**, 912 (1980).

[6] M. Fukugita and T. Yanagida, Phys. Lett. B **174**, 45 (1986).

[7] R. N. Mohapatra, Phys. Rev. **D 34**, 3457 (1986); A. Font, L. Ibanez and F. Quevedo, Phys. Lett **B 228**, 79 (1989): S. P. Martin, Phys. Rev. **D46**, 2769 (1992).

[8] V. A. Kuzmin, JETP Lett. **12**, 228 (1970).

[9] S. L. Glashow, Cargese Lectures (1979).

[10] R. N. Mohapatra and R. E. Marshak, Phys. Rev. Lett. **44**, 1316 (1980).

[11] C. B. Dover, A. Gal and J. M. Richard, Phys. Lett. B **344**, 433 (1995).

[12] W. M. Alberico, A. De Pace and M. Pignone, Nucl. Phys. A **523**, 488 (1991); J. Hufner and B. Z. Kopeliovich, Mod. Phys. Lett. A **13**, 2385 (1998).

[13] E. Friedman and A. Gal, Phys. Rev. D **78**, 016002 (2008).

[14] J. C. Pati and A. Salam, Phys. Rev. D **10**, 275 (1974).

[15] M. Baldo-Ceolin *et al.*, Z. Phys. C **63**, 409 (1994).

[16] M. Takita *et al.* [KAMIOKANDE Collaboration], Phys. Rev. D **34**, 902 (1986); J. Chung *et al.*, Phys. Rev. D **66**, 032004 (2002) [hep-ex/0205093]. For the most recent limit from Super-K, see K. Genezer, Proceedings of the International Workshop on "B-L Violation", LBL (2007); http://inpa.lbl.gov/BLNV/blnv.htm.

[17] Y. A. Kamyshkov, hep-ex/0211006.

[18] A. Young, Proceedings of the International Workshop on "B-L Violation", LBL (2007); http://inpa.lbl.gov/BLNV/blnv.htm.

[19] W. M. Snow, Proceedings of the International Workshop on "B-L Violation", LBL (2007); http://inpa.lbl.gov/BLNV/blnv.htm.

[20] M. Bergevin, Proceedings of the International Workshop on "B-L Violation", LBL (2007); http://inpa.lbl.gov/BLNV/blnv.htm.

[21] S. Rao and R. Shrock, Phys. Lett. B **116**, 238 (1982); J. Pasupathy, Phys. Lett. B **114**, 172 (1982); Riazuddin, Phys. Rev. D **25**, 885 (1982); S. P. Misra and U. Sarkar, Phys. Rev. D **28**, 249 (1983); S. Fajfer and R. J. Oakes, Phys. Lett. B **132**, 433 (1983).

[22] K. S. Babu, P. S. B. Dev and R. N. Mohapatra, Phys. Rev. **D 79**, 015017 (2009); arXiv:0811.3411 [hep-ph].

[23] K. S. Babu, R. N. Mohapatra and S. Nasri, Phys.Rev.Lett. **97**, 131301 (2006).

[24] B. Dutta, Y. Mimura and R. N. Mohapatra, Phys. Rev. Lett. **96**, 061801 (2006)

[25] Z. Chacko and R. N. Mohapatra, Phys. Rev. D **59**, 055004 (1999) [hep-ph/9802388]. V. A. Kuzmin, V. A. Rubakov and M. E. Shaposhnikov, Phys. Lett. B **155**, 36 (1985).

[26] R. N. Mohapatra, N. Okada and H. B. Yu, Phys. Rev. D **77**, 011701 (2008); C. R. Chen *et al.*, W. Klemm, V. Rentala and K. Wang, hep-ph/0811.2105.

[27] N. Arkani-Hamed, S. Dimopoulos and G. R. Dvali, Phys. Lett. B **429**, 263 (1998); I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos and G. R. Dvali, Phys. Lett. B **436**, 257 (1998).

[28] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 3370 (1999)

[29] S. Nussinov and R. Shrock, Phys. Rev. Lett. **88**, 171601 (2002)

[30] N. Arkani-Hamed and M. Schmaltz, Phys. Rev. D **61**, 033005 (2000) [arXiv:hep-ph/9903417].

[31] G. R. Dvali and G. Gabadadze, Phys. Lett. B **460**, 47 (1999)

[32] F. Zwirner, Phys. Lett. B **132** (1983) 103; J. F. Nieves and P. B. Pal, Phys. Lett. B **146**, 406 (1984); R. N. Mohapatra and J. W. F. Valle, Phys. Rev. D **34**, 1642 (1986); J. L. Goity and M. Sher, Phys. Lett. B **346**, 69 (1995) [Erratum-ibid. B **385**, 500 (1996)]; K. S. Babu and R. N. Mohapatra, Phys. Lett. B **518**, 269 (2001).

[33] Goity and Sher, Ref.[32].

[34] G. Costa, F. Feruglio and F. Zwirner, Nuovo Cim. A **70**, 201 (1982); A. Y. Smirnov and F. Vissani, Phys. Lett. B **380**, 317 (1996).

[35] Y. Kamyshkov, private communications.

[36] Z. Berezhiani and L. Bento, Phys. Rev. Lett. **96**, 081801 (2006); for an experimental overview, see talks by Ulrich Schmidt, Andreas Knecht and Anatoly Serebrov, at the " Proceedings of the International Workshop on "B-L Violation", LBL (2007); http://inpa.lbl.gov/BLNV/blnv.htm.